




World Health
Organization

Guideline:
**Calcium supplementation
in pregnant women**



WHO Library Cataloguing-in-Publication Data

Guideline: Calcium supplementation in pregnant women.

1.Calcium, Dietary - administration and dosage. 2.Pregnancy complications, Cardiovascular – prevention and control. 3.Hypertension – prevention and control. 4.Pre-eclampsia – prevention and control. 5.Prenatal nutrition. 6.Dietary supplements. 7.Guideline. I.World Health Organization.

ISBN 978 92 4 150537 6

(NLM classification: QU 130)

© **World Health Organization 2013**

All rights reserved. Publications of the World Health Organization are available on the WHO web site (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press through the WHO web site (http://www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Front Cover picture: Calcium Cristal provided by Quantum MicroMet, Buxton, UK, with their kindly agreement of use.

Design and layout by Full Global Concept S.A.

Suggested citation

WHO. *Guideline: Calcium supplementation in pregnant women*. Geneva, World Health Organization, 2013.

Contents	Acknowledgements	IV
	Financial support	IV
	Executive summary	1
	Scope and purpose	4
	Background	4
	Summary of evidence	5
	Recommendation	7
	Remarks.....	8
	Implications for future research	9
	Dissemination, adaptation and implementation.....	9
	Dissemination	
	Adaptation and implementation	
	Monitoring and evaluation of guideline implementation	
	Guideline development process	10
	Advisory groups	
	Scope of the guideline, evidence appraisal and decision-making	
	Management of conflicts of interest	12
	Plans for updating the guideline.....	13
	References.....	14
Annex 1	Additional analyses	17
Annex 2	GRADE “Summary of findings” tables.....	18
Annex 3	Summary of the Nutrition Guidance Advisory Group’s considerations for determining the strength of the recommendation	20
Annex 4	WHO Steering Committee for Nutrition Guidelines Development	21
Annex 5	Nutrition Guidance Advisory Group – Micronutrients 2010–2011, WHO Secretariat and external resource experts	22
Annex 6	External Experts and Stakeholders Panel.....	27
Annex 7	Questions in Population, Intervention, Control, Outcomes (PICO) format	30

Acknowledgements

This guideline was coordinated by Dr Luz Maria De-Regil under the supervision of Dr Juan Pablo Peña-Rosas, with technical input from Dr João Paulo Dias de Souza, Dr Metin Gulmezoglu, Dr Jose Martines, Dr Matthews Mathai and Dr Lisa Rogers. Thanks are due to the World Health Organization (WHO) Guidelines Review Committee Secretariat staff for their support throughout the process. Thanks are also due to Mr Issa T. Matta and Mrs Chantal Streijffert Garon from the WHO Office of the Legal Counsel for their support in the management of conflicts of interest procedures. Ms Grace Rob and Mrs Paule Pillard from the Evidence and Programme Guidance Unit, Department of Nutrition for Health and Development, provided logistic support.

WHO gratefully acknowledges the technical input of the members of the WHO Steering Committee for Nutrition Guidelines Development and the Nutrition Guidance Advisory Group, especially the chairs of the meetings concerning this guideline, Dr Rafael Flores-Ayala and Dr Emorn Wasantwisut. WHO is also grateful to the Cochrane Pregnancy and Childbirth Group staff for their support in developing and updating the systematic reviews used to inform this guideline.

Financial support

WHO thanks the Government of Luxembourg for financial support for this work.

Guideline¹ Calcium supplementation in pregnant women

Executive summary

Purpose of the guideline: Poor maternal and newborn health and nutrition remain significant contributors to the burden of disease and mortality. Calcium supplementation has the potential to reduce adverse gestational outcomes, in particular by decreasing the risk of developing hypertensive disorders during pregnancy, which are associated with a significant number of maternal deaths and considerable risk of preterm birth, the leading cause of early neonatal and infant mortality. Member States have requested guidance from the World Health Organization (WHO) on the efficacy and safety of calcium supplementation in pregnant women as a public health strategy, in support of their efforts to achieve the Millennium Development Goals and the global targets set in the maternal, infant and child nutrition comprehensive implementation plan. The guideline is intended for a wide audience including policy-makers, their expert advisers, and technical and programme staff at organizations involved in the design, implementation and scaling-up of nutrition actions for public health.

Guideline development methodology: WHO developed the present evidence-informed recommendations using the procedures outlined in the [WHO handbook for guideline development](#). The steps in this process included: (i) identification of priority questions and outcomes; (ii) retrieval of the evidence; (iii) assessment and synthesis of the evidence; (iv) formulation of recommendations, including research priorities; and (v) planning for dissemination, implementation, impact evaluation and updating of the guideline. The *Grading of Recommendations Assessment, Development and Evaluation (GRADE)* methodology was followed to prepare evidence profiles related to preselected topics, based on up-to-date systematic reviews. The guideline development group for nutrition interventions, the Nutrition Guidance Advisory Group, consisted of content experts, methodologists, representatives of stakeholders, consumers and guideline users. These experts participated in two WHO technical consultations concerning this guideline, held in 2011 in Geneva, Switzerland, and Washington DC, United States of America. Members of the External Experts and Stakeholders Panel were identified through a public call for comments, and this panel was involved throughout the guideline development process.

Available evidence: Two Cochrane systematic reviews investigated whether calcium supplementation on a daily basis during pregnancy safely improved maternal and infant outcomes. The findings revealed that this intervention significantly reduced the risk of pre-eclampsia and high blood pressure (with or without proteinuria). Women who received calcium supplements had a significantly higher risk of developing HELLP (haemolysis, elevated liver enzymes, and low platelet count) syndrome, a rare adverse event associated with severe pre-eclampsia. Calcium supplementation had no effects on the risk of developing eclampsia or maternal death or maternal admission to the intensive care unit.

¹ A WHO guideline is any document, whatever its title, containing WHO recommendations about health interventions, whether they be clinical, public health or policy interventions. A recommendation provides information about what policy-makers, health-care providers or patients should do. It implies a choice between different interventions that have an impact on health and that have ramifications for the use of resources. All publications containing WHO recommendations are approved by the WHO Guidelines Review Committee.

In regard to infant outcomes, there was no effect of calcium supplementation on preterm birth (born before 37 weeks' gestation) overall. However, a subgroup analysis suggested that there were fewer preterm births among pregnant women who received between 1.5 g and 2 g of elemental calcium per day than among those women with a lower calcium intake. Calcium supplementation did not have a detectable effect on the risk of low birth weight, admission to a neonatal intensive care unit, stillbirth and neonatal death before hospital discharge.

Recommendation

- In populations where calcium intake is low, calcium supplementation as part of the antenatal care is recommended for the prevention of pre-eclampsia in pregnant women, particularly among those at higher risk of developing hypertension (*strong recommendation*)¹.

A suggested scheme for supplementation in pregnant women is presented in Table 1.

Table 1

Suggested scheme for calcium supplementation in pregnant women

Dosage	1.5–2.0 g elemental calcium/day ^a
Frequency	Daily, with the total daily dosage divided into three doses (preferably taken at mealtimes)
Duration	From 20 weeks' gestation until the end of pregnancy
Target group	All pregnant women, particularly those at higher risk of gestational hypertension ^b
Settings	Areas with low calcium intake

^a 1 g of elemental calcium equals 2.5 g of calcium carbonate or 4 g of calcium citrate.

^b Women are regarded as being at high risk of developing gestational hypertension and pre-eclampsia if they have one or more of the following risk factors: obesity, previous pre-eclampsia, diabetes, chronic hypertension, renal disease, autoimmune disease, nulliparity, advanced maternal age, adolescent pregnancy and conditions leading to hyperplacentation and large placentas (e.g. twin pregnancy). This is not an exhaustive list, but can be adapted/complemented based on the local epidemiology of pre-eclampsia.

Remarks:

- Assessment of the risk of developing gestational hypertensive disorders is to be conducted by a clinician. The clinical management of women with pre-eclampsia or eclampsia requires consideration of other evidence-informed interventions.

¹ A *strong recommendation* is one for which the guideline development group is confident that the desirable effects of adherence outweigh the undesirable effects. Implications of a strong recommendation for patients are that most people in their situation would desire the recommended course of action and only a small proportion would not. Implications for clinicians are that most patients should receive the recommended course of action, and adherence to this recommendation is a reasonable measure of good-quality care. With regard to policy-makers, a strong recommendation means that it can be adapted as a policy in most situations, and for funding agencies it means the intervention likely represents an appropriate allocation of resources (i.e. large net benefits relative to alternative allocation of resources).

-
- Implementation of this recommendation requires close monitoring of women's total daily calcium intake (diet, supplements and antacids). The overall intake of calcium per day should not exceed the locally established upper tolerable limit. In the absence of such reference standards, an upper limit of calcium intake of 3 g/day can be used.
 - The mechanisms through which calcium reduces the risk of gestational hypertension need further elucidation. Available evidence supports the theory that calcium supplementation may reduce the risk of developing pre-eclampsia by filling a dietary gap in calcium intake. In populations where consumption of calcium on average meets the recommended dietary calcium intake, either through calcium-rich foods or fortified staple foods, calcium supplementation is not encouraged as it may not improve the outcomes related to pre-eclampsia and hypertensive disorders of pregnancy but might increase the risk of adverse effects. Although antacids are a rich source of calcium, they are not part of the diet and their use should be limited to the treatment of heartburn or indigestion. The calcium content of any other vitamin and mineral supplements that are also being taken should be considered when recommending calcium supplementation, to reduce the risk of hypercalcaemia.
 - Determination of the dietary calcium intake of an individual woman is a complex task. The target group for this recommendation comprises populations with observed low dietary calcium intake or those living in geographical areas where calcium-rich foods are not commonly available or consumed. Calcium intake at population level can be estimated through various means including dietary surveys using 24-hour recalls, food frequency questionnaires or food weighing, as well as through secondary data estimates derived from the Food and Agriculture Organization (FAO) food balance sheets or household consumption and expenditure surveys.
 - Healthy dietary practices to promote adequate calcium intake through local calcium-rich foods should be encouraged in the general population, including pregnant women.
 - Interaction between iron supplements and calcium supplements may occur, although the consequences of prolonged calcium supplementation for iron status among different age groups are still unclear. Therefore, the two nutrients should preferably be administered several hours apart (i.e. iron may be consumed between meals) rather than concomitantly.
 - Selection of the most appropriate delivery platform should be context specific, with the aim of reaching the most vulnerable populations and ensuring a timely and continuous supply of supplements. Calcium supplementation could be delivered by lay health workers along with targeted monitoring and evaluation.
 - Calcium supplements are available as tablets or capsules. Tablets (soluble tablets, effervescent tablets, chewable tablets for use in the mouth and modified-release tablets) are solid dosage forms containing one or more active ingredients.

Research priorities: Guideline group members and stakeholders identified several research priorities to improve the body of evidence at the basic, clinical, epidemiological and operational levels on both benefits and harms of this intervention among pregnant women. They are listed in the guideline.

Scope and purpose

This guideline provides global, evidence-informed recommendations on calcium supplementation as a public health intervention for the purpose of improving maternal and infant health outcomes.

The guideline will help Member States and their partners in their efforts to make informed decisions on the appropriate nutrition actions to achieve the Millennium Development Goals, in particular, reduction of child mortality (MDG 4) and improvement of maternal health (MDG 5). It will also support Member States in their efforts to achieve global targets on the maternal, infant and young child nutrition comprehensive implementation plan, especially global target 3, which entails achieving a relative reduction of 30% in the number of infants born with a weight <2500 g by the year 2025 (1). The guideline is intended for a wide audience including policy-makers, their expert advisers, and technical and programme staff at organizations involved in the design, implementation and escalation of nutrition actions in public health programmes.

This document presents the key recommendation and a summary of the supporting evidence. Further details of the evidence base are provided in Annexes 1 and 2 and other documents listed in the references.

Background

Poor maternal and newborn health and nutrition remain significant contributors to the burden of disease. In 2010, 3.1 million babies died in the first 28 days of life, mostly due to low birth weight, severe infections, asphyxia and preterm birth. Every year, 15 million babies are born prematurely, of whom 1.1 million die in the neonatal period or in infancy. In addition, many of those who survive have a lifetime disability such as learning disabilities and/or visual and hearing problems (2).

Approximately 287 000 women died during pregnancy and childbirth in 2010, mostly due to maternal health complications (3). Hypertensive disorders of pregnancy include (pre-existing) chronic hypertension and gestational hypertension, pre-eclampsia and eclampsia (4,5). These disorders complicate approximately 2-8% of all pregnancies and have been associated with preterm and low birth weight and maternal mortality (4). Pre-eclampsia is diagnosed when gestational hypertension (maternal blood pressure $\geq 140/90$ mmHg for the first time in the second half of pregnancy) is accompanied by proteinuria >300 mg in a 24-hour period. The pathogenesis of pre-eclampsia has not been thoroughly elucidated, however, it is related to disturbances in placentation in early pregnancy, followed by generalized inflammation and progressive endothelial damage. Pre-eclampsia can be classified as mild or severe. In severe pre-eclampsia, the blood pressure is $\geq 160/110$ mmHg and there is proteinuria ≥ 2 g/24 h, with or without substantial maternal organ damage (5). Such end-organ damage as a result of pre-eclampsia can present with haemolysis, elevated liver enzymes and low platelet count, a constellation of symptoms known as HELLP syndrome (5). This is a rare condition that occurs in 10–20% of cases with severe pre-eclampsia (6).

Calcium is the most abundant mineral in the body and is essential for many diverse processes, including bone formation, muscle contraction, and enzyme and hormone functioning (7). Most of the body's calcium is found

in the bones and teeth; approximately 1% is present in the intracellular structures, cell membrane and extracellular fluids (8). Calcium absorption increases during pregnancy and no additional intake is needed (9). A dietary intake of 1200 mg/day of calcium for pregnant women is recommended by WHO and the Food and Agriculture Organization of the United Nations (FAO) (7). Inadequate consumption of this nutrient by pregnant women can lead to adverse effects in both the mother and the fetus, including osteopenia, tremor, paraesthesia, muscle cramping, tetanus, delayed fetal growth, low birth weight and poor fetal mineralization (10).

Serum calcium concentrations are maintained within narrow limits in the body and thus have limited use for the assessment of calcium nutritional status at both the individual and the population levels. Calcium intake could be a useful indicator of status at the population level. The main dietary sources of this nutrient are milk, dairy products, calcium-set tofu and fortified foods (9); some local foods such as lime-treated corn meal also have abundant calcium. There is no information on the adequacy of calcium intake worldwide. However, some studies at the regional and national levels suggest that low calcium intake at population level occurs frequently (11, 12).

Various studies have suggested that calcium supplementation during pregnancy has a beneficial effect on reducing the risk of pregnancy-induced hypertension (10). The results of trials evaluating the effect of supplementation on maternal bone mineral density, fetal mineralization, and preterm birth, however, are less conclusive (13). Excessive consumption of calcium may increase the risk of urinary stones and urinary tract infection, and reduce the absorption of other essential micronutrients (13).

In supplements, calcium is present in the form of carbonate, citrate, lactate or gluconate, and in general all these forms have good bioavailability (14, 15). At least one salt of calcium for oral administration (in a variety of doses) is included in most national essential medicines lists (16), calcium carbonate being the most common. As calcium carbonate has the highest content of elemental calcium (40%), it may have the best efficacy-cost ratio in pregnancy (17), but this needs to be confirmed in future analyses.

Summary of evidence

An existing Cochrane systematic review was updated (13) and another review was newly developed (18) to specifically investigate whether daily calcium supplementation in pregnancy safely improves maternal and infant outcomes. The maternal outcomes considered critical for decision-making by the Nutrition Guidance Advisory Group were pre-eclampsia, eclampsia, high blood pressure with or without proteinuria, complications at delivery, and any adverse events. Important infant outcomes were preterm birth, low birth weight (<2500 g), stillbirth, death during the neonatal period, and any adverse effects such as being small for gestational age or admission to a neonatal care unit.

For the purposes of this guideline, the results of the two systematic reviews were combined, thus including the findings of 21 randomized controlled trials (RCT), involving more than 19 000 pregnant women from both developed and developing countries in all continents. These trials compared calcium supplementation with receiving a placebo or no

intervention in addition to regular antenatal care. The total daily dosage of supplemental calcium ranged between 300 mg (0.3 g) and 2000 mg (2 g). Most of the studies started supplementation at 20 weeks' gestation and were considered of high quality.

For all women, irrespective of the baseline risk of developing hypertension and calcium intake status, calcium supplementation more than halved the risk of pre-eclampsia when compared with a placebo (average risk ratio (RR) 0.48, 95% confidence interval (CI) 0.34–0.67, 15 trials, 16 490 women). The risk reduction was 41% in women at low risk of developing hypertension (RR 0.59, 95% CI 0.42–0.82, 10 trials, 15 903 women) whereas among those at high risk of hypertensive disorders, a much higher risk reduction of 78% was recorded (RR 0.22, 95% CI 0.12–0.42, five trials, 587 women). Although the women's response to calcium supplementation was heterogeneous in terms of the magnitude of the effect, there was a consistent protective effect of the intervention (Annex 1, Figure 1; Annex 2).

High blood pressure (with or without proteinuria) showed, in general, a similar pattern to that of pre-eclampsia. Overall, fewer women on calcium supplementation had high blood pressure compared with those receiving placebo (RR 0.65, 95% CI 0.53–0.81, 12 trials, 15 470 women). Women at high risk of developing hypertension showed the greatest reduction in risk (RR 0.47, 95% CI 0.22–0.90, four trials, 327 women), closely followed by those with a low baseline dietary calcium intake (RR 0.44, 95% CI 0.28–0.70, seven trials, 10 418 women).

With regard to the outcome of eclampsia, there were no significant differences between women on calcium supplementation and women receiving placebo or no treatment (RR 0.66, 95% CI 0.40–1.11, five trials, 14 185 women). Similar effects were found in a non-Cochrane review analysis restricted to developing countries (19).

A significant increase in the risk ratio for HELLP (haemolysis, elevated liver enzymes, low platelet counts) syndrome was observed in women who received calcium supplementation compared with those who received placebo (RR 2.67, 95% CI 1.05–6.82, two trials, 12 901 women). There were no effects on maternal death (RR 0.17, 95% CI 0.02–1.39, one trial, 8312 women) or maternal admission to the intensive care unit (RR 0.84, 95% CI 0.66–1.07, one trial, 8312 women).

In regard to infant outcomes (Annex 2), there was no effect of calcium supplementation on preterm birth, although a subgroup analysis by supplemental dose suggested that among pregnant women consuming ≥ 1.5 g of calcium per day fewer babies were born before 37 weeks' gestation than among those receiving < 1.5 g per day (RR 0.78, 95% CI 0.63–0.98 vs. RR 0.72, 95% CI 0.08–6.52 (Annex 1, Figure 2)). There were no significant differences between women who received calcium supplements and those who did not in the risk of having a low-birth-weight baby (RR 0.85, 95% CI 0.72–1.01, nine trials, 14 883 infants) or in the risk of the baby being admitted to neonatal intensive care unit (RR 1.05, 95% CI 0.94–1.18, four trials, 14 062 women) and, stillbirth or neonatal death before hospital discharge (RR 0.90, 95% CI 0.74–1.09, 11 trials, 15 665 women). One of the systematic reviews assessed additional infant outcomes (18) and noted that a trial conducted in 1978 with 273 women reported that calcium supplementation might increase infant bone mineral density. There was no evidence of an effect of calcium

supplementation on intrauterine growth restriction, birth length or fetal head circumference.

The quality of the evidence for admission to a neonatal intensive care unit was high, while it was moderate for pre-eclampsia, high blood pressure, and maternal admission to an intensive care unit. The quality of the evidence was low for eclampsia, maternal death, HELLP (haemolysis, elevated liver enzymes, low platelet counts) syndrome, preterm birth, low birth weight and perinatal mortality.

Recommendation

- In populations where calcium intake is low, calcium supplementation as part of the antenatal care is recommended for the prevention of pre-eclampsia among pregnant women, particularly among those at higher risk of hypertension (*strong recommendation*)^{1,2}.

A suggested scheme for supplementation in pregnant women is presented in Table 1.

Table 1

Suggested scheme for calcium supplementation in pregnant women

Dosage	1.5–2.0 g elemental calcium/day ^a
Frequency	Daily, with the total daily dosage divided into three doses (preferably taken at mealtimes)
Duration	From 20 weeks' gestation until the end of pregnancy
Target group	All pregnant women, particularly those at higher risk of gestational hypertension ^b
Settings	Areas with low calcium intake

^a 1 g of elemental calcium equals 2.5 g of calcium carbonate or 4 g of calcium citrate.

^b Women are regarded as being at high risk of developing hypertension and pre-eclampsia if they have one or more of the following risk factors: obesity, previous pre-eclampsia, diabetes, chronic hypertension, renal disease, autoimmune disease, nulliparity, advanced maternal age, adolescent pregnancy and conditions leading to hyperplacentation and large placentas (e.g. twin pregnancy). This is not an exhaustive list, but can be adapted/complemented based on the local epidemiology of pre-eclampsia.

¹ A *strong recommendation* is one for which the guideline development group is confident that the desirable effects of adherence outweigh the undesirable effects. Implications of a strong recommendation for patients are that most people in their situation would desire the recommended course of action and only a small proportion would not. Implications for clinicians are that most patients should receive the recommended course of action, and adherence to this recommendation is a reasonable measure of good-quality care. With regard to policy-makers, a strong recommendation means that it can be adapted as a policy in most situations, and for funding agencies it means the intervention likely represents an appropriate allocation of resources (i.e. large net benefits relative to alternative allocation of resources).

² Considerations of the guideline development group for determining the strength of the recommendation are summarized in Annex 3.

- Remarks**
- Assessment of the risk of developing gestational hypertensive disorders is to be conducted by a clinician. The clinical management of women with pre-eclampsia or eclampsia requires consideration of other evidence-informed interventions (5).
 - Implementation of this recommendation requires close monitoring of women's total daily calcium intake (diet, supplements and antacids). The overall intake of calcium per day should not exceed the locally established upper tolerable limit. In the absence of such reference standards, an upper limit of calcium intake of 3 g/day can be used (7).
 - The mechanisms through which calcium reduces the risk of gestational hypertension need further elucidation. Available evidence supports the theory that calcium supplementation may reduce the risk of developing pre-eclampsia by filling a dietary gap in calcium intake (5). In populations where consumption of calcium on average meets the recommended dietary calcium intake, either through calcium-rich foods or fortified staple foods, calcium supplementation is not encouraged as it may not improve the outcomes related to pre-eclampsia and hypertensive disorders of pregnancy but might increase the risk of adverse effects. Although antacids are not a rich source of calcium, they are not part of the diet and their use should be limited to the treatment of heartburn or indigestion. The calcium content of any other vitamin and mineral supplements that are also being taken should be considered when recommending calcium supplementation, to reduce the risk of hypercalcaemia.
 - Determination of the dietary calcium intake of an individual woman is a complex task. The target group for this recommendation comprises populations with observed low dietary calcium intake or those living in geographical areas where calcium-rich foods are not commonly available or consumed (5). Calcium intake at population level can be estimated through various means including dietary surveys using 24-hour recalls, food frequency questionnaires or food weighing, as well as through secondary data estimates derived from FAO food balance sheets or household consumption and expenditure surveys (20, 21).
 - Healthy dietary practices to promote adequate calcium intake through local calcium-rich foods should be encouraged in the general population, including pregnant women (5).
 - Interaction between iron supplements and calcium supplements may occur, although the consequences of prolonged calcium supplementation for iron status among different age groups are still unclear (22–25). Therefore, the two nutrients should preferably be administered several hours apart (i.e. iron may be consumed between meals) rather than concomitantly.
 - Selection of the most appropriate delivery platform should be context-specific, with the aim of reaching the most vulnerable populations and ensuring a timely and continuous supply of supplements. Calcium supplementation could be delivered by lay health workers along with targeted monitoring and evaluation (26).
 - Calcium supplements are available as tablets or capsules. Tablets (soluble tablets, effervescent tablets, chewable tablets for use in the mouth and modified-release tablets) are solid dosage forms containing one or more active ingredients (27).

Implications for future research

Discussion with members of the Nutrition Guidance Advisory Group and stakeholders highlighted the limited evidence available in some areas, meriting further research on calcium supplementation in pregnant women, in particular, in the following areas:

- biological mechanisms underlying the relationships between calcium supplementation, pre-eclampsia and HELLP (haemolysis, elevated liver enzymes, low platelet counts) syndrome;
- minimal dose and optimal commencement of supplementation to achieve a positive effect on pre-eclampsia and other gestational outcomes such as preterm birth;
- effects of calcium supplementation on maternal and infant outcomes in conjunction with other nutrients (e.g. vitamin D) or as part of a supplement with multiple vitamins and minerals;
- effects of gestational calcium supplementation among adolescents;
- long-term effects of calcium supplementation during pregnancy on mother and infant health;
- additional benefit of calcium supplementation among pregnant women who are already receiving antihypertensive treatment; and
- operational research assessing delivery mechanisms, compliance, acceptability and costs of providing separate versus multiple micronutrients. Calcium supplementation programmes need careful monitoring and evaluation to assess their successes and failures in terms of integration into the overall antenatal care package.

Dissemination, adaptation and implementation

Dissemination

The current guideline will be disseminated through electronic media such as slide presentations, CD-ROMs and the World Wide Web, through the WHO Nutrition mailing list (into which the Micronutrients mailing list was merged) and United Nations Standing Committee on Nutrition (SCN) mailing list, social media, the [WHO nutrition web site](#), and the [WHO e-Library of Evidence for Nutrition Actions](#) (eLENA). The WHO e-Library of Evidence for Nutrition Actions compiles and displays WHO guidelines related to nutrition, along with complementary documents such as systematic reviews and other evidence that informed the guidelines, biological and behavioural rationales, and additional resources produced by Member States and global partners. In addition, the guideline will be disseminated through a broad network of international partners, including WHO country and regional offices, ministries of health, WHO collaborating centres, universities, other United Nations agencies and nongovernmental organizations. It will also be published in and disseminated via the [WHO Reproductive Health Library](#).

Adaptation and implementation

As this is a global guideline it should be adapted to the context of each Member State. Prior to implementation, a public health programme that includes the provision of calcium supplements to pregnant women should have well-defined objectives that take into account available resources, existing policies, suitable delivery platforms and suppliers, communication

channels, and potential stakeholders. Ideally, calcium supplementation should be implemented as part of an integrated programme for antenatal care and preceded by an assessment of the calcium intake at population level. The latter can be estimated through various means including dietary surveys using 24-hour recalls, food frequency questionnaires or food weighing, as well as through secondary data estimates derived from FAO food balance sheets or household consumption and expenditure surveys (20, 21).

To ensure that WHO global guidelines and other evidence-informed recommendations for nutrition interventions are better implemented in low and middle-income countries, the Department of Nutrition for Health and Development works with the WHO Evidence-Informed Policy Network (EVIPNet) programme. EVIPNet promotes partnerships at country level between policy-makers, researchers and civil society to facilitate policy development and implementation through use of the best available evidence.

Monitoring and evaluation of guideline implementation

A plan for monitoring and evaluation with appropriate indicators is encouraged at all stages. The impact of this guideline can be evaluated within countries (i.e. monitoring and evaluation of the programmes implemented at national or regional scale) and across countries (i.e. the adoption and adaptation of the guideline globally). The WHO Department of Nutrition for Health and Development, Evidence and Programme Guidance Unit, jointly with the Centers for Disease Control and Prevention (CDC) International Micronutrient Malnutrition Prevention and Control (IMMPaCt) programme, and with input from international partners, has developed a generic logic model for micronutrient interventions in public health (28) to depict the plausible relationships between inputs and expected MDGs by applying the micronutrient programme evaluation theory. Member States can adjust the model and use it in combination with appropriate indicators, for designing, implementing, monitoring and evaluating the successful escalation of nutrition actions in public health programmes.

For evaluation at the global level, the WHO Department of Nutrition for Health and Development has developed a centralized platform for sharing information on nutrition actions in public health practice implemented around the world. By sharing programmatic details, specific country adaptations and lessons learnt, this platform will provide examples of how guidelines are being translated into nutrition actions.

Guideline development process

This guideline was developed in accordance with WHO evidence-informed guideline development procedures, as outlined in the [WHO handbook for guideline development](#) (29).

Advisory groups

The WHO Steering Committee for Nutrition Guidelines Development (Annex 4), led by the Department of Nutrition for Health and Development, was established in 2009 with representatives from all WHO departments with an interest in the provision of scientific nutrition advice, including Maternal, Neonatal, Child and Adolescent Health and Development and Reproductive Health and Research. The WHO Steering Committee for Nutrition Guidelines Development meets twice yearly and both guided and provided overall

supervision of the guideline development process. Two additional groups were formed: a guideline development group and a panel of external experts and stakeholders.

The Nutrition Guidance Advisory Group (Annex 5) was established in 2009. A subgroup for micronutrients was established for the biennium 2010–2011. Its role was to advise WHO on the choice of important outcomes for decision-making and in the interpretation of the evidence. The Nutrition Guidance Advisory Group includes experts from various [WHO expert advisory panels](#) and those identified through open calls for specialists, taking into consideration a balanced gender mix, multiple disciplinary areas of expertise and representation from all WHO regions. Efforts were made to include content experts, methodologists, representatives of potential stakeholders (such as managers and other health professionals involved in the health-care process) and consumers. Representatives of commercial organizations may not be members of a WHO guideline development group.

The External Experts and Stakeholders Panel (Annex 6) was consulted on the scope of the guideline, the questions addressed and the choice of important outcomes for decision-making, as well as with regard to review of the completed draft guideline. This was done through the WHO Micronutrients and [SCN](#) mailing lists that together included over 5500 subscribers, and through the [WHO nutrition web site](#).

Scope of the guideline, evidence appraisal and decision-making

An initial set of questions (and the components of the questions) to be addressed in the guideline was the critical starting point for formulating the recommendation. The questions were drafted by technical staff at the Evidence and Programme Guidance Unit, Department of Nutrition for Health and Development, based on policy and programme guidance needs of Member States and their partners. The population, intervention, control, outcomes (PICO) format was used (Annex 7). The questions were discussed and reviewed by the WHO Steering Committee for Nutrition Guidelines Development, and feedback was received from four stakeholders.

A Nutrition Guidance Advisory Group meeting was held on 14–16 March 2011 in Geneva, Switzerland, to finalize the scope of the questions and rank the critical outcomes and populations of interest for the recommendation on calcium supplementation in pregnant women for the improvement of maternal and neonatal outcomes. The Nutrition Guidance Advisory Group – Micronutrients Subgroup discussed the relevance of the questions and modified them as needed. The guideline group scored the relative importance of each outcome from 1 to 9 (where 7–9 indicated that the outcome was critical for a decision, 4–6 indicated that it was important and 1–3 indicated that it was not important). The final key questions on this intervention, along with the outcomes that were identified as critical for decision-making, are listed in PICO format in Annex 7.

Two systematic reviews (13, 18) were used to summarize and appraise the evidence using the Cochrane methodology for randomized controlled

trials¹. WHO staff prepared evidence summaries according to the *Grading of Recommendations Assessment, Development and Evaluation (GRADE)* approach to assess the overall quality of the evidence (30). GRADE considers: the study design; the limitations of the studies in terms of their conduct and analysis; the consistency of the results across the available studies; the directness (or applicability and external validity) of the evidence with respect to the populations, interventions and settings where the proposed intervention may be used; and the precision of the summary estimate of the effect.

Both the systematic reviews and the GRADE evidence profiles for each of the critical outcomes were used for drafting this guideline. The draft recommendation was discussed by the WHO Steering Committee for Nutrition Guidelines Development and at a second consultation with the Nutrition Guidance Advisory Group, held on 7–9 November 2011 in Washington DC, United States of America. At the second consultation, the guideline development group members independently voted on the strength of the recommendation, taking into account: (i) the desirable and undesirable effects of the intervention; (ii) the quality of the available evidence; (iii) values and preferences related to the intervention in different settings; and (iv) the cost of options available to health-care workers in different settings (Annex 3). It was noted how they balanced the decisions for each of the four domains. The voting results and the summary of the considerations for establishing the strength of the recommendation were disclosed before the end of the meeting and further discussed as needed. Consensus was defined as agreement by simple majority of the guideline group members. WHO staff present at the meeting as well as other external technical experts involved in the collection and grading of the evidence were not allowed to vote. There were no strong disagreements among the guideline group members.

A public call for comments on the final draft guideline was released in 2012. All interested stakeholders became members of the External Experts and Stakeholders Panel but were allowed to comment on the draft guideline only after submitting a signed Declaration of Interests form. Feedback was received from 46 stakeholders. WHO staff addressed each comment and then finalized the guideline and submitted it for clearance by WHO before publication.

This guideline complements the report of the WHO technical consultation on the prevention and treatment of pre-eclampsia and eclampsia, held on 7–8 April 2011 in Geneva, Switzerland (5).

Management of conflicts of interest

According to the rules in the WHO *Basic documents* (31), all experts participating in WHO meetings must declare any interest relevant to the meeting prior to their participation. The conflicts of interest statements for all guideline group members were reviewed by the responsible technical officer and the relevant departments before finalization of the group composition

¹ The detailed methods used in each systematic review, as well as their search dates, are published and available (open access) via The Cochrane Library. As part of the Cochrane pre-publication editorial process, this review was commented on by external peers (an editor, and two referees external to the editorial team) and the group's statistical adviser (<http://www.cochrane.org/cochrane-reviews>). The *Cochrane handbook for systematic reviews of interventions* describes in detail the process of preparing and maintaining Cochrane systematic reviews on the effects of health-care interventions.

and invitation to attend a guideline group meeting. All guideline group members and participants of the guideline development meetings submitted a Declaration of Interests form along with their curriculum vitae before each meeting. In addition, they verbally declared potential conflicts of interest at the beginning of each meeting. The procedures for management of conflicts of interest strictly followed WHO *Guidelines for declaration of interests (WHO experts)* (32). The potential conflicts of interest declared by the members of the guideline group are summarized below.

- Dr Héctor Bourges Rodriguez declared being chair of the executive board of the Danone Institute in Mexico, a non-profit organization promoting research and dissemination of scientific knowledge in nutrition, and receiving funds as chair honorarium from this organization. Some activities of the Danone Institute in Mexico may generally relate to nutrition and are funded by Danone Mexico, a food producer.
- Dr Emorn Wasantwisut declared serving as a technical/scientific adviser to the International Life Sciences Institute (ILSI)/South East Asia's Food and Nutrients in Health and Disease Cluster and as a reviewer of technical documents and speaker for Mead Johnson Nutritionals. Her research unit received funds for research support from Sight and Life and the International Atomic Energy Agency for the use of stable isotopes to define interactions of vitamin A and iron.

Plans for updating the guideline

This guideline will be reviewed in 2016. If new information is available at that time, a guideline review group will be convened to evaluate the new evidence and revise the recommendation if needed. The Department of Nutrition for Health and Development at the WHO headquarters in Geneva, along with its internal partners, will be responsible for coordinating the guideline update, following formal [WHO handbook for guideline development](#) (29) procedures. WHO welcomes suggestions regarding additional questions for evaluation when the guideline is due for review.

References

1. Resolution WHA65.11. Nutrition. Maternal, infant and young child nutrition: draft comprehensive implementation plan. In: *Sixth-fifth World Health Assembly, Geneva, 21–26 May 2012. Resolutions and decisions, and list of participants*. Geneva, World Health Organization, 2012 (A65/11) Annex: 5–23 (http://apps.who.int/gb/ebwha/pdf_files/WHA65/A65_11-en.pdf, accessed 13 June 2013).
2. March of Dimes, PMNCH, Save the Children, WHO. *Born too soon: the Global Action Report on Preterm Birth*. Howson CP, Kinney MV, Lawn JE, eds. Geneva, World Health Organization, 2012.
3. WHO. *World Health Statistics 2013*. Geneva, World Health Organization, 2013 (<http://apps.who.int/iris/bitstream/10665/81965/1/9789241564588-eng.pdf>, accessed 13 June 2013).
4. Duley L. The global impact of pre-eclampsia and eclampsia. *Seminars in perinatology* 2009, 33(3):130-7.
5. *WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia*. Geneva, World Health Organization, 2011 (http://whqlibdoc.who.int/publications/2011/9789241548335_eng.pdf, accessed 25 March 2013).
6. Haram K, Svendsen E, Abildgaard U. The HELLP syndrome: clinical issues and management. A Review. *BMC Pregnancy Childbirth* 2009, 9:8 (<http://www.biomedcentral.com/1471-2393/9/8/>, accessed 13 June 2013).
7. World Health Organization, Food and Agricultural Organization of the United Nations. *Vitamin and mineral requirements in human nutrition*, 2nd ed. Geneva, World Health Organization, 2004 (<http://www.who.int/nutrition/publications/micronutrients/9241546123/en/index.html>, accessed 13 June 2013).
8. Sauberlich HE. *Laboratory tests for the assessment of nutritional status*, 2nd ed. Boca Raton, CRC Press, 1999.
9. National Institute of Medicine. *Dietary reference intakes for Calcium and vitamin D*. New York, National Academies Press, 2011.
10. Villar J et al. Methodological and technical issues related to the diagnosis, screening, prevention and treatment of pre-eclampsia and eclampsia. *International Journal of Gynecology and Obstetrics*, 2004, 85(Suppl. 1):S28–S41.
11. Gibson RS, Cavalli-Sforza T. Using reference nutrient density goals with food balance sheet data to identify likely micronutrient deficits for fortification planning in countries in the Western Pacific region. *Food and Nutrition Bulletin*, 2012, 33(Suppl. 3):S214–220.
12. Broadley MR et al. Dietary requirements for magnesium, but not calcium, are likely to be met in Malawi based on National Food Supply Data. *International Journal of Vitamin and Nutrition Research*, 2012, 82(3):192–199.
13. Hofmeyr GJ et al. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database of Systematic Reviews*, 2010, (8):CD001059.

14. Heaney RP et al. Absorbability and cost effectiveness in calcium supplementation. *Journal of the American College of Nutrition*, 2001, 20(3):239–246.
15. Straub DA. Calcium supplementation in clinical practice: a review of forms, doses, and indications. *Nutrition Clinical Practice*, 2007, 22:286.
16. *National Medicines List/Formulary/Standard Treatment Guidelines*. Geneva, World Health Organization, 2012 [cited 2012 November 1st] (http://www.who.int/selection_medicines/country_lists/en/index.html, accessed 13 June 2013).
17. Management Sciences for Health. *International drug price indicator guide 2011 edition*. Arlington, Management Sciences for Health Inc, 2012 (http://erc.msh.org/dmpguide/pdf/DrugPriceGuide_2011_en.pdf, accessed 13 June 2013).
18. Buppasiri P et al. Calcium supplementation (other than for preventing or treating hypertension) for improving pregnancy and infant outcomes. *Cochrane Database of Systematic Reviews*, 2011, (10):CD007079.
19. Imdad A, Jabeen A, Bhutta ZA. Role of calcium supplementation during pregnancy in reducing risk of developing gestational hypertensive disorders: a meta-analysis of studies from developing countries. *BMC Public Health*, 2011, 11(Suppl. 3):S18.
20. Dary O, Imhoff-Kunsch B. Measurement of food consumption to inform food fortification and other nutrition programs: an introduction to methods and their application. *Food and Nutrition Bulletin*, 2012, 33(Suppl. 3):S141–145.
21. Fiedler JL et al. Household Consumption and Expenditures Surveys (HCES): a primer for food and nutrition analysts in low- and middle-income countries. *Food and Nutrition Bulletin*, 2012, 33(Suppl. 3):S170–184.
22. Kalkwarf HJ, Harrast SD. Effects of calcium supplementation and lactation on iron status. *American Journal of Clinical Nutrition*, 1998, 67(6):1244–1249.
23. Mølgaard C, Kaestel P, Michaelsen KF. Long-term calcium supplementation does not affect the iron status of 12-14-y-old girls. *American Journal of Clinical Nutrition*, 2005, 82(1):98–102.
24. Bendich A. Calcium supplementation and iron status of females. *Nutrition*, 2001, 17(1):46–51.
25. Gaitán D et al. Calcium does not inhibit the absorption of 5 milligrams of nonheme or heme iron at doses less than 800 milligrams in nonpregnant women. *Journal of Nutrition*, 2011, 141(9):1652–1656.
26. *WHO recommendations. Optimizing health worker roles to improve access to key maternal and newborn health interventions through task shifting*. Geneva, World Health Organization, 2012 (http://apps.who.int/iris/bitstream/10665/77764/1/9789241504843_eng.pdf, accessed 25 March 2013).

-
27. *The international pharmacopoeia*, 4th ed. Volumes 1 and 2. Geneva, World Health Organization, 2008 (<http://apps.who.int/phint/en/p/about/>, accessed 13 June 2013).
 28. WHO/CDC. *Logic model for micronutrient interventions in public health*. Vitamin and Mineral Nutrition Information System. Geneva, World Health Organization, 2011 (WHO/NMH/NHD/MNM/11.5; http://www.who.int/vmnis/toolkit/logic_model/en/index.html, accessed 13 June 2013).
 29. *WHO handbook for guideline development*. Guidelines Review Committee. Geneva, World Health Organization, 2012. (http://apps.who.int/iris/bitstream/10665/75146/1/9789241548441_eng.pdf, accessed 25 March 2013).
 30. Guyatt G et al. GRADE guidelines 1. Introduction – GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology*, 2011, 64:383–394.
 31. *Basic documents*, 47th ed. Geneva, World Health Organization, 2009 (<http://apps.who.int/gb/bd/>, accessed 13 June 2013).
 32. *Guidelines for declaration of interests (WHO experts)*. Geneva, World Health Organization, 2010.

Annex 1. Additional analyses

Figure 1. Effect of gestational calcium supplementation on pre-eclampsia in comparison with receiving a placebo or no intervention, by risk of developing hypertension





Figure 2. Effect of gestational calcium supplementation on preterm birth in comparison with receiving a placebo or no intervention, by calcium supplementation dose



* Boggess (1997) reported on premature labour
For details of studies, see reference (13, 18).

Annex 2. GRADE “Summary of findings” tables

Calcium supplementation for pregnant women: maternal outcomes

Patient or population: Pregnant women

Settings: All settings

Intervention: Calcium supplementation

Comparison: Placebo or no intervention

Outcomes	Relative effect or mean difference (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)*	Comments
Pre-eclampsia	RR 0.48 (0.34–0.67)	16 490 (15 trials)	⊕⊕⊕⊖ moderate ¹	
Eclampsia	RR 0.66 (0.40–1.11)	14 185 (5 trials)	⊕⊕⊖⊖ low ^{1,2}	
High blood pressure (with or without proteinuria) ⁺	RR 0.65 (0.53–0.81)	15 470 (12 trials)	⊕⊕⊕⊖ moderate ¹	
Maternal death ^{+♣}	RR 0.17 (0.02–1.39)	8312 (1 trial)	⊕⊕⊖⊖ low ^{2,3}	
Maternal admission to intensive care unit ^{+♣}	RR 0.84 (0.66–1.07)	8312 (1 trial)	⊕⊕⊕⊖ moderate ³	
Haemolysis, elevated liver enzymes, low platelet counts (HELLP) syndrome ⁺	RR 2.67 (1.05–6.82)	12 901 (2 trials)	⊕⊕⊖⊖ low ^{2,4}	

CI, confidence interval; RR, risk ratio.

*GRADE Working Group grades of evidence:

High quality: We are very confident that the true effect lies close to that of the estimate of the effect.

Moderate quality: We have moderate confidence in the effect estimate. The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low quality: Our confidence in the effect estimate is limited. The true effect may be substantially different from the estimate of the effect.

Very low quality: We have very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of the effect.

¹ Serious statistical heterogeneity possibly due to variation in baseline dietary intake of calcium, but there is consistency in the direction of the effect.

² Wide confidence intervals (imprecision).

³ Only one study reported on this outcome.

⁴ Few events.

+ For details of studies included in the review, see reference (13).

♣ For details of studies included in the review, see reference (18).

Calcium supplementation for pregnant women: newborn outcomes

Patient or population: Pregnant women

Settings: All settings

Intervention: Calcium supplementation

Comparison: Placebo or no intervention

Outcomes	Relative effect or mean difference (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)*	Comments
Preterm birth	RR 0.83 (0.66–1.04)	16 093 (14 trials)	⊕⊕⊖⊖ low ^{1,2}	
Low birth weight ⁺	RR 0.85 (0.72–1.01)	14 883 (9 trials)	⊕⊕⊖⊖ low ^{1,2}	
Perinatal mortality ⁺	RR 0.84 (0.61–1.16)	5145 (7 trials)	⊕⊕⊖⊖ low ^{1,2}	
Admission to neonatal intensive care unit ^{+♣}	RR 1.05 (0.94–1.18)	14 062 (4 trials)	⊕⊕⊕⊕ high	

CI, confidence interval; RR, risk ratio.

*GRADE Working Group grades of evidence:

High quality: We are very confident that the true effect lies close to that of the estimate of the effect.

Moderate quality: We have moderate confidence in the effect estimate. The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low quality: Our confidence in the effect estimate is limited. The true effect may be substantially different from the estimate of the effect.

Very low quality: We have very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of the effect.

¹ Serious statistical heterogeneity possibly explained by the supplemental dose of calcium, but there is consistency in the direction of the effect.

² There are some trials at high risk of bias, particularly due to high losses to follow-up and lack of blinding.

+ For details of studies included in the review, see reference (13).

♣ For details of studies included in the review, see reference (18).

Annex 3. Summary of the Nutrition Guidance Advisory Group’s considerations for determining the strength of the recommendation

<p>Quality of evidence:</p>	<ul style="list-style-type: none"> • Low and moderate-quality evidence for most outcomes • Clear impact on pre-eclampsia and possibly on preterm birth • Low-quality evidence for adverse effects
<p>Values and preferences:</p>	<ul style="list-style-type: none"> • Effects greater in those with low calcium intake and larger doses of calcium • Evidence is consistent for pre-eclampsia and preterm birth, both of which are responsible for a considerable proportion of the burden of maternal and infant morbidity and mortality • Some members noted that the two largest studies do not show a clinical impact of this intervention
<p>Trade-off between benefits and harms:</p>	<ul style="list-style-type: none"> • Current evidence suggests that benefits outweigh disadvantages, particularly in populations with low calcium intake • The possibility of developing HELLP syndrome and renal stones is a concern and more evidence is needed for this outcome
<p>Costs and feasibility:</p>	<ul style="list-style-type: none"> • This intervention may be adopted as a policy when adequate health systems and delivery platforms already exist • There is a lack of information on the cost of this intervention • This intervention may require intensive resources to target those women with an increased risk of hypertension

Annex 4. WHO Steering Committee for Nutrition Guidelines Development

Dr Ala Alwan

Acting Director
Department of Chronic Diseases
and Health Promotion
Noncommunicable Diseases and
Mental Health (NMH) Cluster

Dr Francesco Branca

Director
Department of Nutrition for Health
and Development
Noncommunicable Diseases and
Mental Health (NMH) Cluster

Dr Ruediger Krech

Director
Department of Ethics, Equity, Trade
and Human Rights
Information, Evidence and
Research (IER) Cluster

Dr Knut Lonnroth

Medical Officer
The Stop TB Strategy
HIV/AIDS, TB and Neglected
Tropical Diseases (HTM) Cluster

Dr Daniel Eduardo Lopez Acuña

Director
Department of Strategy, Policy and
Resource Management
Health Action in Crises (HAC)
Cluster

Dr Elizabeth Mason

Director
Department of Maternal, Neonatal,
Child and Adolescent Health and
Development
Family and Community Health
(FCH) Cluster

Dr Michael Mbizvo

Director
Department of Reproductive
Health and Research
Family and Community Health
(FCH) Cluster

Dr Jean-Marie Okwo-Bele

Director
Department of Immunization,
Vaccines and Biologicals
Family and Community Health
(FCH) Cluster

Dr Gottfried Otto Hirschall

Director
Department of HIV/AIDS
HIV/AIDS, TB and Neglected
Tropical Diseases (HTM) Cluster

Dr Isabelle Romieu

Director
Dietary Exposure Assessment
Group, Nutrition and Metabolism
Section
International Agency for Research
on Cancer (IARC), Lyons, France

Dr Sergio Spinaci

Associate Director
Global Malaria Programme
HIV/AIDS, TB and Neglected
Tropical Diseases (HTM) Cluster

Dr Willem Van Lerberghe

Director
Department of Health Policy,
Development and Services
Health Systems and Services (HSS)
Cluster

Dr Maged Younes

Director
Department of Food Safety,
Zoonoses and Foodborne Diseases
Health Security and Environment
(HSE) Cluster

Dr Nevio Zagaria

Acting Director
Department of Emergency
Response and Recovery
Operations
Health Action in Crises (HAC)
Cluster

Annex 5. Nutrition Guidance Advisory Group Micronutrients 2010–2011, WHO Secretariat and external resource experts

A. Nutrition Guidance Advisory Group – Micronutrients

(Note: the areas of expertise of each guideline group member are given in italics)

Ms Deena Alasfoor

Ministry of Health
Muscat, Oman
*Health programme management,
food legislations,
surveillance in primary health care*

Dr Beverley-Ann Biggs

International and Immigrant
Health Group
Department of Medicine
University of Melbourne
Parkville, Australia
*Micronutrients supplementation,
clinical infectious diseases*

Dr Héctor Bourges Rodríguez

Instituto Nacional de Ciencias
Medicas y Nutrición Salvador
Zubiran
Mexico City, Mexico
*Nutritional biochemistry and
metabolism research,
food programmes, policy, and
regulations*

Dr Norm Campbell

Departments of Medicine
Community Health Sciences and
Physiology and Pharmacology
University of Calgary
Calgary, Canada
*Physiology and pharmacology,
hypertension
prevention and control*

Dr Rafael Flores-Ayala

Centers for Disease Control and
Prevention (CDC)
Atlanta, United States of America
*Nutrition and human capital
formation, nutrition and
growth, impact of micronutrient
interventions*

Professor Malik Goonewardene

Department of Obstetrics and
Gynaecology
University of Ruhuna
Galle, Sri Lanka
*Obstetrics and gynaecology, clinical
practice*

Dr Junsheng Huo

National Institute for Nutrition and
Food Safety
Chinese Center for Disease Control
and Prevention
Beijing, China
*Food fortification, food science and
technology,
standards and legislation*

Dr Janet C. King

Children's Hospital Oakland
Research Institute
Oakland, United States of America
*Micronutrients, maternal and child
nutrition,
dietary requirements*

Dr Marzia Lazzerini

Department of Paediatrics and
Unit of Research on Health
Services and International Health
Institute for Maternal and Child
Health
IRCCS Burlo Garofolo
Trieste, Italy
*Paediatrics, malnutrition, infectious
diseases*

Professor Malcolm E. Molyneux

College of Medicine – University of
Malawi
Blantyre, Malawi
*Malaria, international tropical
diseases research and practice*

Engineer Wisam Qarqash
Jordan Health Communication
Partnership
Johns Hopkins University
Bloomberg School of Public Health
Amman, Jordan
*Design, implementation and
evaluation of health
communications and programmes*

Dr Daniel Raiten
Office of Prevention Research and
International Programs
National Institutes of Health (NIH)
Bethesda, United States of America
*Malaria, maternal and child health,
human development research*

Dr Mahdi Ramsan Mohamed
Research Triangle Institute (RTI)
International
Dar es Salaam, the United Republic
of Tanzania
*Malaria control and prevention,
neglected tropical diseases*

Dr Meera Shekar
Health Nutrition Population
Human Development Network
(HDNHE)
The World Bank
Washington, DC, United States of
America
*Costing of interventions in public
health nutrition,
programme implementation*

Dr Rebecca Joyce Stoltzfus
Division of Nutritional Sciences
Cornell University
Ithaca, United States of America
*International nutrition and public
health, iron and vitamin A nutrition,
programme research*

Ms Carol Tom
East, Central and Southern Africa
Health Community (ECSA-HC)
Nairobi, Kenya
*Food fortification technical
regulations and standards, policy
harmonization*

Dr David Tovey
The Cochrane Library
Cochrane Editorial Unit
London, England
*Systematic reviews, health
communications,
evidence for primary health care*

Mrs Vilma Qahoush Tyler
UNICEF Regional Office for Central
and Eastern Europe
and the Commonwealth of
Independent States (CEE/CIS)
Geneva, Switzerland
*Food fortification, public health
programmes*

Dr Gunn Elisabeth Vist
Department of Preventive and
International Health
Norwegian Knowledge Centre for
the Health Services
Oslo, Norway
*Systematic review methods and
evidence assessment using GRADE
methodology*

Dr Emorn Wasantwisut
Mahidol University
Nakhon Pathom, Thailand
*International nutrition,
micronutrient biochemistry
and metabolism*

B. WHO Secretariat

Mr Daniel Albrecht
Technical Officer
Social Determinants of Health Unit
Department of Ethics, Equity, Trade
and Human Rights Information

Dr Maria del Carmen Casanovas
Technical Officer
Evidence and Programme
Guidance Unit
Department of Nutrition for Health
and Development

Dr Bernadette Daelmans

Medical Officer
Newborn and Child Health and
Development Unit
Department of Maternal, Newborn,
Child and Adolescent Health and
Development

Dr Luz Maria De-Regil

Epidemiologist
Evidence and Programme
Guidance Unit
Department of Nutrition for Health
and Development

Dr Chris Duncombe

Medical Officer
Anti-retroviral Treatment and HIV
Care Unit
Department of HIV/AIDS

Dr Olivier Fontaine

Medical Officer
Newborn and Child Health and
Development Unit
Department of Maternal, Newborn,
Child and Adolescent Health and
Development

Dr Ahmet Metin Gulmezoglu

Medical Officer
Technical Cooperation with
Countries for Sexual and
Reproductive Health
Department of Reproductive
Health and Research

Dr José Martines

Coordinator
Newborn and Child Health and
Development Unit
Department of Maternal, Newborn,
Child and Adolescent Health and
Development

Dr Matthews Mathai

Medical Officer Department
of Maternal, Newborn, Child
and Adolescent Health and
Development

Dr Mario Merialdi

Coordinator
Improving Maternal and Perinatal
Health Unit
Department of Reproductive
Health and Research

Dr Juan Pablo Peña-Rosas

Coordinator
Evidence and Programme
Guidance Unit Department
of Nutrition for Health and
Development

Dr Lisa Rogers

Technical Officer
Evidence and Programme
Guidance Unit
Department of Nutrition for Health
and Development

Dr João Paulo Dias de Souza

Medical Officer
Technical Cooperation with
Countries for Sexual and
Reproductive Health
Department of Reproductive
Health and Research

Dr Severin Von Xylander

Medical Officer
Department of Maternal, Newborn,
Child and Adolescent Health and
Development

Dr Godfrey Xuereb

Technical Officer
Surveillance and Population-based
Prevention Unit
Department of Chronic Diseases
and Health Promotion

C. WHO regional offices

Dr Abel Dushimimana

Medical Officer
Nutrition
WHO Regional Office for Africa
Brazzaville, Congo

Dr Evelina Chapman

Specialist
Health Research Management
WHO Regional Office for the
Americas/Pan American Health
Organization
Washington, DC, United States of
America

Dr Luis Gabriel Cuervo

Senior Advisor
Research Promotion and
Development
WHO Regional Office for the
Americas/Pan American Health
Organization
Washington, DC, United States of
America

Dr Ruben Grajeda

Micronutrients Technical Advisor
Child and Adolescent Health
WHO Regional Office for the
Americas/Pan American Health
Organization
Washington, DC, United States of
America

Dr Chessa Lutter

Regional Adviser
Child and Adolescent Health
WHO Regional Office for the
Americas/Pan American Health
Organization
Washington, DC, United States of
America

Dr Gina Tambini

Area Manager
Family and Community Health
WHO Regional Office for the
Americas/Pan American Health
Organization
Washington, DC, United States of
America

Dr Kunal Bagchi

Regional Adviser
Nutrition and Food Safety
WHO Regional Office for South-
East Asia
New Delhi, India

Dr Joao Breda

Noncommunicable Diseases and
Environment
WHO Regional Office for Europe
Copenhagen, Denmark

Mrs Trudy Wijnhoven

Technical Officer, Nutrition
Surveillance
Noncommunicable Diseases and
Environment
WHO Regional Office for Europe
Copenhagen, Denmark

Dr Ayoub Al-Jawaldeh

Regional Adviser
Nutrition
WHO Regional Office for the
Eastern Mediterranean
Cairo, Egypt

Dr Tommaso Cavalli-Sforza

Regional Adviser
Nutrition
WHO Regional Office for the
Western Pacific
Manila, Philippines

D. External resource experts

Dr Denise Coitinho Delmuè

United Nations System Standing
Committee on Nutrition (SCN)
Geneva, Switzerland

Dr Lynnette Neufeld

Micronutrient Initiative
Ottawa, Canada

Dr. Lely Solari Zerpa

Centro Nacional de Salud Pública
Instituto Nacional de Salud
Lima, Peru

Mr Arnold Timmer

United Nations Children's Fund
(UNICEF)
New York, United States of America

Dr Carol L Wagner

Medical University of South
Carolina,
Charleston, United States of
America

Dr Connie M. Weaver

Purdue University
West Lafayette, United States of
America

Annex 6. External Experts and Stakeholders Panel

Professor Leila Alouane

National Institute of Nutrition and Food Technology (INNTA)
Tunis, Tunisia

Dr Stephanie Atkinson

McMaster University
Hamilton, Canada

Dr Salmeh Bahmanpour

Shiraz University of Medical Sciences
Shiraz, Iran (Islamic Republic of)

Mr Eduard Baladia Rodríguez

Spanish Association of Dietitians and Nutritionists
Barcelona, Spain

Mr Julio Basulto Marset

Spanish Association of Dietitians and Nutritionists
Barcelona, Spain

Professor Hiba Bawadi

Jordan University of Science and Technology
Irbid, Jordan

Ms Laura Begueria

Association of Enterprises of Dietetic Products and Food Supplements (AFEPADI)
Barcelona, Spain

Ms Cristiana Berti

North-West University
Vanderbijlpark, South Africa

Mr Ashok Bhurtyal

World Health Organization,
Country Office for Nepal
Kathmandu, Nepal

Professor Steven Boyages

University of Sydney
Sydney, Australia

Professor Cyrus Cooper

International Osteoporosis Foundation
Nyon, Switzerland

Dr Rosa Corcoy

Hospital de la Santa Creu i Sant Pau
Barcelona, Spain

Dr Anne Davis

University of Saint Joseph
West Hartford, United States of America

Mr David Eboku

Uganda National Bureau of Standards
Nakawa, Uganda

Dr Mario Flores

National Institute of Public Health
Cuernavaca, Mexico

Ms Rae Galloway

PATH
Seattle, United States of America

Dr Frederick Grant

International Potato Center
Nairobi, Kenya

Mr Rufus Greenbaum

Stanmore, United Kingdom of Great Britain and Northern Ireland

Dr Suzanne Harris

International Life Sciences Institute
Washington DC, United States of America

Dr Nicholas Harvey

International Osteoporosis Foundation
Nyon, Switzerland

Dr Nilmini Hemachandra

Family Health Bureau, Ministry of Health
Colombo, Sri Lanka

Dr Valerie Holmes

Queen's University Belfast
Belfast, United Kingdom of Great
Britain and Northern Ireland

Dr Elina Hypponen

University College London
Institute of Child Health
London, United Kingdom of Great
Britain and Northern Ireland

Dr Yustina Anie Indriastuti

Indonesian Medical Association
Physician Nutrition
Jakarta, Indonesia

Dr Bharati Kulkarni

National Institute of Nutrition
Indian Council of Medical Research
Hyderabad, India

Dr Maria Manera

Spanish Association of Dietitians-
Nutritionists
Barcelona, Spain

Professor Nayeli Macias Morales

National Institute of Public Health
Cuernavaca, Mexico

Dr Dominic Moyo

Ministry of Health
Lilongwe, Malawi

Dr Maria Mulhern

University of Ulster
Coleraine, United Kingdom of
Great Britain and Northern Ireland

Dr Madhavan K Nair

National Institute of Nutrition
(Indian Council of Medical
Research)
Hyderabad, India

Ms Victoria Ojo

University of Ibadan
Oyo State, Nigeria

Dr Moshood Omotayo

Cornell University
Ithaca, United States of America

Dr Kirsty Pourshahidi

University of Ulster
Coleraine, United Kingdom of
Great Britain and Northern Ireland

Ms Dawna Royall

Dietitians of Canada
Fergus, Canada

Mrs Laurence Rycken

International Dairy Federation
Brussels, Belgium

Dr Jaganath Sharma

John Snow Inc.
Boston, United States of America

Dr Dina Shehab

National Nutrition Institute
Cairo, Egypt

Mr Fred Tabung

University of South Carolina
Columbia, United States of
America

Dr Rob te Biesebeke

Nestlé
Vevey, Switzerland

Mr Andrew Thorne-Lyman

Harvard School of Public Health
Boston, United States of America

Dr Thach S. Tran

Australian Research Centre for
Health of Women and Babies
University of Adelaide
Adelaide, Australia

Dr Tserendolgor Uush

Nutrition Research Center
Public Health Institute
Ulaanbaatar, Mongolia

Dr Napaphan Viriyautsahakul

Ministry of Public Health
Nonthaburi, Thailand



Dr Mostafa Waly

Sultan Qaboos University
Muscat, Oman

Dr Rianne Weggemans

Health Council of the Netherlands
The Hague, the Netherlands

Mr You Zhou

Minerva Foundation Institute for
Medical Research
Helsinki, Finland

Annex 7. Questions in Population, Intervention, Control, Outcomes (PICO) format

Effects of calcium supplementation in women during pregnancy

- a. Should calcium supplements be given to pregnant women for improvement of maternal and infant health outcomes?
- b. If so, at what dose, frequency and duration of the intervention?

Population:	<ul style="list-style-type: none"> • Pregnant women • Subpopulations (listed in order of priority): <ul style="list-style-type: none"> – Populations with a low versus adequate baseline dietary calcium intake – Populations with above average risk versus low or average risk of hypertensive disorders of pregnancy
Intervention:	<ul style="list-style-type: none"> • Oral calcium supplements • Oral calcium supplements given in combination with other micronutrients • Subgroup analyses: <ul style="list-style-type: none"> – By dose of calcium – By calcium compound: calcium carbonate, lactate, gluconate – By regimen: daily versus other – By duration of supplementation – By trimester of pregnancy in which supplementation was started
Control:	<ul style="list-style-type: none"> • Placebo or no treatment • Micronutrient supplements without calcium (to assess the additive effect of calcium)
Outcomes:	<p>Maternal</p> <ul style="list-style-type: none"> • High blood pressure with or without proteinuria • High blood pressure with significant proteinuria (pre-eclampsia) • Eclampsia (the occurrence of one or more convulsions (fits) in association with pre-eclampsia) • Complications at delivery (assisted delivery) • Bone softness • Osteoporosis • Any adverse effects <p>Infant</p> <ul style="list-style-type: none"> • Low birth weight (<2500 g) • Birth weight • Preterm birth (<37 weeks' gestation) • Length at birth • Admission to a neonatal intensive care unit • Stillbirth or death in early neonatal period (0–7 days of life) • Any adverse effects • Small for gestational age
Setting:	All settings/global

For more information, please contact:

Department of Nutrition for Health and Development
World Health Organization
Avenue Appia 20, CH-1211 Geneva 27, Switzerland
Fax: +41 22 791 4156
E-mail: nutrition@who.int
www.who.int/nutrition



ISBN 978 92 4 150537 6

